

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 30-1136: Anti-CD56 / NCAM Monoclonal Antibody (Clone:MEM-188)

Clonality: Monoclonal **Clone Name:** MEM-188 Application: FACS Reactivity: Human Gene: NCAM1 Gene ID: 4684 **Uniprot ID:** P13591 Format: **Purified Alternative Name:** NCAM1,NCAM Isotype: Mouse IgG2a

Immunogen Information: KG-1 human acute myelogenous leukemia cell line

## **Description**

CD56 (NCAM, neural cell adhesion molecule) is a transmembrane glycoprotein of immunoglobulin family serving as adhesive molecule which is ubiquitously expressed in nervous system, usually as 120 kDa, 140 kDa or 180 kDa isoform, and it is also found on T cells and NK cells. Polysialic modification results in reduction of CD56-mediated cell adhesion and is involved in cell migration, axonal growth, pathfinding and synaptic plasticity. CD56 is a widely used neuroendocrine marker with a high sensitivity for neuroendocrine tumours and ovarian granulosa cell tumours.

## **Product Info**

Amount: 0.1 mg

**Purification :** Purified by protein-A affinity chromatography

**Storage condition :** Store at 2-8°C. Do not freeze.

## **Application Note**

Flow Cytometry Recommended dilution:3 µg/ml

Immunoprecipitation Immunohistochemistry Recommended dilution: 1:50

Pretreatment: Heat retrieval (microwave oven) in citrate buffer, standard ABC visualization.

Positive tissue: Positive membrane-specific reaction in adjacent peripheral nerve

Staining of paraprostatic autonomic ganglion.

Staining technique: Standard ABC technique (DAB+).