

30-2028: Anti-CD79a Monoclonal Antibody (Clone:HM57)-FITC Conjugated

| | |
|--------------------------------|--|
| Clonality : | Monoclonal |
| Clone Name : | HM57 |
| Application : | FACS, IHC, IHC-Fr |
| Reactivity : | Human, Pig, Mouse, Rat, Bovine, Equine, Guinea pig, Opossum, Rabbit, Chicken |
| Conjugate : | FITC |
| Gene : | CD79A |
| Gene ID : | 973 |
| Uniprot ID : | P11912 |
| Alternative Name : | CD79A,IGA,MB1 |
| Isotype : | Mouse IgG1 |
| Immunogen Information : | Synthetic peptide corresponding to amino acids 202-216 of human CD79a |

Description

CD79a (Ig alpha, MB1) forms disulfide-linked heterodimer with CD79b (Ig beta). They both are transmembrane proteins with extended cytoplasmic domains containing immunoreceptor tyrosine activation motives (ITAMs), and together with cell surface immunoglobulin they constitute B-cell antigen-specific receptor (BCR). CD79a and b are the first components of BCR that are expressed developmentally. They appear on pro-B cells in association with the endoplasmic reticulum chaperone calnexin. Subsequently, in pre-B cells, CD79 heterodimer is associated with lambda5-VpreB surrogate immunoglobulin and later with antigen-specific surface immunoglobulins. At the plasma cell stage, CD79a is present as an intracellular component. CD79a/b complex interacts with Src-family tyrosine kinase Lyn, which phosphorylates its cytoplasmic ITAM motives to form docking sites for downstream signaling.

Product Info

| | |
|----------------------------|---|
| Amount : | 100 tests |
| Storage condition : | Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. |




Figure 1: Intracellular staining of CD79a in human peripheral blood with anti-CD79a (HM57) FITC.