

## 30-2089: Anti-CD79b Monoclonal Antibody (Clone:CB3-1)-FITC Conjugated

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	CB3-1
<b>Application :</b>	FACS, IP, WB
<b>Reactivity :</b>	Human
<b>Conjugate :</b>	FITC
<b>Gene :</b>	CD79B
<b>Gene ID :</b>	974
<b>Uniprot ID :</b>	P40259
<b>Alternative Name :</b>	CD79B,B29,IGB
<b>Isotype :</b>	Mouse IgG1
<b>Immunogen Information :</b>	Fraction of Ig-associated molecules isolated from Ramos B cells

### Description

CD79b (Ig beta, B29) forms disulfide-linked heterodimer with CD79a (Ig alpha, MB1). They both are transmembrane proteins with extended cytoplasmic domains containing immunoreceptor tyrosine activation motives (ITAMs), and together with cell surface immunoglobulin they constitute B-cell antigen-specific receptor (BCR). CD79a and b are the first components of BCR that are expressed developmentally. They appear on pro-B cells in association with the endoplasmic reticulum chaperone calnexin. Subsequently, in pre-B cells, CD79 heterodimer is associated with lambda5-VpreB surrogate immunoglobulin and later with antigen-specific surface immunoglobulins. CD79a/b complex interacts with Src-family tyrosine kinase Lyn, which phosphorylates its cytoplasmic ITAM motives to form docking sites for downstream signaling.

### Product Info

<b>Amount :</b>	100 tests
<b>Storage condition :</b>	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.




Figure 1: Surface staining of CD79b in human peripheral blood with anti-CD79b (CB3-1) FITC.