

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-1408: Anti-CD79b Monoclonal Antibody (Clone:HM79)(Discontinued)

Clonality: Monoclonal Clone Name: HM79

Application: FACS, IP, WB, ICC

Reactivity: Mouse
Gene: Cd79b
Gene ID: 15985
Uniprot ID: P15530
Format: Purified
Alternative Name: Cd79b,lgb
Isotype: Hamster lgG2

Immunogen Information: Purified CD79a/b (alpha/beta) dimers from WEHI231 cells

Description

CD79b (Ig beta, B29) forms disulfide-linked heterodimer with CD79a (Ig alpha, MB1). They both are transmembrane proteins with extended cytoplasmic domains containing immunoreceptor tyrosine activation motives (ITAMs), and together with cell surface immunoglobulin they constitute B-cell antigen-specific receptor (BCR). CD79a and b are the first components of BCR that are expressed developmentally. They appear on pro-B cells in association with the endoplasmic reticulum chaperone calnexin. Subsequently, in pre-B cells, CD79 heterodimer is associated with lambda5-VpreB surrogate immunoglobulin and later with antigen-specific surface immunoglobulins. CD79a/b complex interacts with Src-family tyrosine kinase Lyn, which phosphorylates its cytoplasmic ITAM motives to form docking sites for downstream signaling.

Product Info

Amount: 0.1 mg

Purification : Purified by protein-A affinity chromatography

Storage condition : Store at 2-8°C. Do not freeze.

Application Note

Flow Cytometry Recommended dilution:1 Ã\[\hat{A}\pug/m\] Immunoprecipitation Western Blotting Immunocytochemistry