

## 30-2421: Anti-CD8 Monoclonal Antibody (Clone:MEM-31)-PerCP Conjugated

| Clonality :           | Monoclonal   |
|-----------------------|--|
| Clone Name :          | MEM-31   |
| Application :         | FACS, IP   |
| Reactivity :          | Human  |
| Conjugate :           | PerCP  |
| Gene :                | CD8A   |
| Gene ID :             | 925  |
| Uniprot ID :          | P01732   |
| Alternative Name :    | T-lymphocyte differentiation antigen T8/Leu-2, CD8a, MAL |
| Isotype :             | Mouse IgG2a  |
| Immunogen Information | : Crude thymus membrane fraction.                        |

## Description

The CD8 T cell coreceptor (monomer approx. 32-34 kDa) is expressed as alpha/beta heterodimer on majority of MHC Irestricted conventional T cells and thymocytes and as alpha/alpha homodimer on subsets of memory T cells, intraepithelial lymphocytes, NK cells and dendritic cells. Regulation of CD8 beta level on T cell surface seems to be an important mechanism to control their effector function. Assembly of CD8 alpha-beta but not alpha-alpha dimers is connected with formation or localization to the lipid rafts. Recruiting triggered TCR complexes to these membrane microdomains as well as affinity of TCR to MHC I is modulated by CD8, thereby affecting the functional diversity of the TCR signaling.

## **Product Info**

Amount :100 testsStorage condition :Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.




Figure 1: Surface staining of human peripheral blood using anti-human CD8 (clone MEM-31) PerCP.