

## 30-1481: Anti-CD83 Monoclonal Antibody (Clone:HB15e)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	HB15e
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human
<b>Gene :</b>	CD83
<b>Gene ID :</b>	9308
<b>Uniprot ID :</b>	Q01151
<b>Format :</b>	Purified
<b>Alternative Name :</b>	CD83
<b>Isotype :</b>	Mouse IgG1
<b>Immunogen Information :</b>	Human CD83-transfected Cos cells

### Description

CD83 is a 40-45 kDa heavily glycosylated type I cell surface glycoprotein of immunoglobulin family. It is expressed on the surface of mature dendritic cells, Langerhans cells in the skin, and interdigitating reticulum cells in the lymphoid tissues. Low expression of CD83 has been reported in activated T and B cells. Cytoplasmic expression of CD83 can be detected also in monocytes and macrophages. CD83 is involved in modulation of antigen presentation. Soluble CD83 has immunoregulatory functions, it is able to down-regulate dendritic cell maturation and stimulation of T cells. In the developing immune system, release of soluble CD83 from dendritic cells upon stimulation by gram-positive or gram-negative bacteria has anti-allergic effect. Herpes simplex virus, on the other hand, causes CD83 degradation in mature dendritic cells.

### Product Info

<b>Amount :</b>	0.1 mg
<b>Purification :</b>	Purified by protein-A affinity chromatography
<b>Storage condition :</b>	Store at 2-8°C. Do not freeze.

### Application Note

**Flow Cytometry** *Positive control:* peripheral blood-derived dendritic cells

**Immunohistochemistry (frozen sections)** *Application note:* acetone fixation




Figure 1: Flow cytometry surface staining pattern of human interferon alpha stimulated peripheral whole blood stained using anti-human CD83 (HB15e) purified antibody (concentration in sample 2 µg/ml, GAM APC).




Figure 2: Separation of human monocytes (red-filled) from CD83 negative lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human interferon alpha stimulated peripheral whole blood stained using anti-human CD83 (HB15e) purified antibody (concentration in sample 2 µg/ml, GAM APC).