

### 30-1238: Anti-CD95 / Fas Monoclonal Antibody (Clone:LT95)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	LT95
<b>Application :</b>	FACS, IHC
<b>Reactivity :</b>	Human
<b>Gene :</b>	FAS
<b>Gene ID :</b>	355
<b>Uniprot ID :</b>	P25445
<b>Format :</b>	Purified
<b>Alternative Name :</b>	Tumor necrosis factor receptor superfamily member 6, Apo-1 antigen, Apoptosis-mediating surface antigen FAS
<b>Isotype :</b>	Mouse IgG1
<b>Immunogen Information :</b>	HUT-78 human T cell lymphoma cell line

#### Description

CD95 (Fas, APO-1), a 46 kDa transmembrane glycoprotein, is a cell death receptor of the TNFR superfamily. Stimulation of CD95 results in aggregation of its intracellular death domains, formation of the death-inducing signaling complex (DISC) and activation of caspases. In type I cells caspase 3 is activated by high amounts of caspase 8 generated at the DISC, in type II cells low concentration of caspase 8 activates pathway leading to the release of cytochrome c from mitochondria and activation of caspase 3 by cytochrome c. Besides its roles in induction of apoptosis, Fas also triggers pro-inflammatory cytokine responses.

#### Product Info

<b>Amount :</b>	0.1 mg
<b>Purification :</b>	Purified by protein-A affinity chromatography
<b>Storage condition :</b>	Store at 2-8°C. Do not freeze.

#### Application Note

**Flow Cytometry** excellent **Immunohistochemistry** *Positive tissue: tonsil*