

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-1919: FITC Conjugated, Anti-Cytokeratin (Pan-reactive) Monoclonal Antibody (Clone:C-11)

Clonality: Monoclonal

Clone Name: C-11
Application: WB
Conjugate: FITC

Isotype: Mouse IgG1

Immunogen Information: Keratin-enriched preparation from human epidermoid carcinoma cell line A431.

Description

Cytokeratins are a subfamily of intermediate filaments and characterized by remarkable biochemical diversity. Cytokeratins are represented in epithelial tissues by at least 20 different polypeptides, molecular weight between 40 kDa and 68 kDa. The individual cytokeratin polypeptides are designated 1 to 20 and divided into the type I (acidic cytokeratins 9-20) and type II (basic to neutral cytokeratins 1-8) families.

Product Info

Amount: 0.1 mg

Storage condition: Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood spiked with HeLa cells stained using anti-Cytokeratins (C-11) FITC antibody (concentration in sample 1.5 μ g/ml).




Figure 2: Anti-Hu Cytokeratins FITC antibody (clone C-11) specificity verification by flow cytometry Anti-Cytokeratins (C-11) FITC antibody (concentration in sample 1.5 μ g/ml, red-filled histogram) binds specifically to intracellular cytokeratins in permeabilized HeLa cells (upper panel), but does not stain permeabilized TF-1 cells (lower panel). Level of non-specific binding was assessed using Mouse IgG1 isotype control FITC antibody (MOPC-21) under same conditions (concentration in sample 1.5 μ g/ml, black-dashed histogram).