

## 36-1556: Monoclonal Antibody to Progesterone Receptor (Marker of Progesterin Dependence)(Clone : PR484)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	PR484
<b>Application :</b>	IHC
<b>Reactivity :</b>	Human
<b>Gene :</b>	PGR
<b>Gene ID :</b>	5241
<b>Uniprot ID :</b>	P06401
<b>Format :</b>	Purified
<b>Alternative Name :</b>	PGR,NR3C3
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Recombinant human Progesterone Receptor protein

### Description


This MAb is specific to progesterone receptor and shows minimal cross-reaction with other members of the family. Progesterone receptor is expressed as two major isoforms, PR-A (81kDa) and PR-B (116kDa). Expression of PgR has been suggested to reflect an intact estrogen regulatory machinery and therefore, predict better clinical response to endocrine therapy than ER alone. It is excellent for immunohistochemical staining of formalin/paraffin tissues.

### Product Info

<b>Amount :</b>	100 µg
<b>Purification :</b>	Affinity Chromatography
<b>Content :</b>	100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
<b>Storage condition :</b>	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),



Formalin-fixed, paraffin-embedded human Breast Carcinoma stained with Progesterone Receptor Ab (PR484).