







Email: info@abeomics.com

30-2205: PE Conjugated Anti-IgG (Fab) Monoclonal Antibody (Clone:4A11)

Clonality: Monoclonal

* abeomics

Clone Name: 4A11 **FACS** Application: Reactivity: Human Conjugate: PΕ Gene: IGHG1 **Uniprot ID:** P01857 **Alternative Name:** IGHG1 Isotype: Mouse IgG2b

Immunogen Information: Purified human IgG.

Description

Immunoglobulin G (IgG) is a 150 kDa soluble protein that serves as a major effector molecule of the humoral immune response in man. Its concentration in blood plasma of healthy individuals is approximately 10 g/l, which accounts for about 75% of the total plasma immunoglobulins. IgG has the highest stability of blood immunoglobulins (T1/2 = 21 days) and is able of placental transfer. IgG is secreted by plasma cells at a comparably high rate as other immunoglobulins.

Product Info

Amount: 0.1 mg

Storage condition: Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human IgG (Fab) (A8B5) PE antibody (concentration in sample 9 μg/ml).




Figure 2: Flow cytometry multicolor surface staining pattern of human lymphocytes stained using anti-human CD19 (LT19) APC antibody (10 µl reagent / 100 µl of peripheral whole blood) and antihuman IgG (Fab) (A8B5) PE antibody (concentration in sample 9 µg/ml).



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982






Figure 3: Separation of human IgG (Fab) positive monocytes (red-filled) from IgG (Fab) negative CD19 negative lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human IgG (Fab) (A8B5) PE antibody (concentration in sample 15 $\mu g/ml$).