

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-1216: Anti-p21Waf1 Monoclonal Antibody (Clone:WA-1)(Discontinued)

Clonality: Monoclonal

Clone Name: WA-1
Application: WB
Reactivity: Human
Gene: CDKN1A
Gene ID: 1026
Uniprot ID: P38936
Format: Purified

Alternative Name: CDKN1A,CAP20,CDKN1,CIP1,MDA6,PIC1,SDI1,WAF1

Isotype: Mouse IgG1

Immunogen Information: Human p21 protein

Description

The tumour suppressor p21Waf1 (Wild-type p53-Activated Fragment 1; also known as Cip1, Cdk interacting protein, or SDI 1) is a cyclin-dependent kinase (Cdk) inhibitor, which is expressed by involvement of p53, Egr-1, AP2, STATs or other transcription factors upon various stimuli resulting in cell cycle arrest. Through its N-terminal domain p21Waf1 inhibits Cdk activity, whereas through the C-terminal domain it inhibits the activity of PCNA (proliferating cell nuclear antigen) to activate DNA replication. Cytosolic location of p21 counteracts its inhibitory activities.

Product Info

Amount: 0.1 mg

Purification : Purified by protein-A affinity chromatography

Storage condition : Store at 2-8°C. Do not freeze.

Application Note

Immunofluoroscence Immunohistochemistry Positive tissue: colon carcinoma




Figure 1: Immunohistochemistry staining of colon carcinoma (paraffin-embedded sections) with anti-p21 (WA-1).



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com



Figure 2: Immunofluorescence staining of p21 in human primary fibroblasts using anti-p21 (WA-1; green). Actin cytoskeleton decorated by phalloidin (red) and cell nuclei stained with DAPI (blue).