

## 30-1506: Anti-SCIMP Monoclonal Antibody (Clone:NVL-07)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	NVL-07
<b>Application :</b>	FACS, IP, WB, ICC
<b>Reactivity :</b>	Human
<b>Gene :</b>	SCIMP
<b>Gene ID :</b>	388325
<b>Uniprot ID :</b>	Q6UWF3
<b>Format :</b>	Purified
<b>Alternative Name :</b>	SCIMP,C17orf87,UNQ5783/PRO16090
<b>Isotype :</b>	Mouse IgG2a
<b>Immunogen Information :</b>	Recombinant intracellular part of human SCIMP

### Description

SCIMP (SLP adaptor and Csk interacting membrane protein), also known as Nvl, is a palmitoylated transmembrane adaptor protein expressed in professional antigen presenting cells, most prominently in the lymph nodes and spleen. It is associated with tetraspanin-enriched microdomains (together with MHC II). There is a close relationship between SCIMP and tyrosinkinase Lyn, which is constitutively bound to it by its SH3 domain. After MHC II-mediated stimulation in the immunological synapse SCIMP becomes phosphorylated at several tyrosine residues and provides docking sites for Grb2 and SLP65 or SLP76 adaptors transducing the signal downstream, as well as for the kinase Csk with modulatory roles.

### Product Info

<b>Amount :</b>	0.1 mg
<b>Purification :</b>	Purified by protein-A affinity chromatography
<b>Storage condition :</b>	Store at 2-8°C. Do not freeze.




Figure 1: Intracellular staining of SCIMP in a population of HEK-293T-SCIMP transfectants using monoclonal antibody (clone NVL-07, purified). The staining pattern reflects heterogeneity in the cell population regarding transfection efficiency.