

30-2286: PE Conjugated Anti-TCR alpha/beta Monoclonal Antibody (Clone:IP26)

Clonality :	Monoclonal
Clone Name :	IP26
Application :	FACS
Reactivity :	Human
Conjugate :	PE
Gene :	PTCRA
Gene ID :	171558
Uniprot ID :	Q6ISU1
Alternative Name :	PTCRA
Isotype :	Mouse IgG1

Description

The antigen-specific T cell receptor (TCR) is composed of either alpha and beta subunit, or gamma and delta subunit. Majority of T cells present in the blood, lymph and secondary lymphoid organs express TCR alpha/beta heterodimers, whereas the T cells expressing TCR gamma/delta heterodimers are localized mainly in epithelial tissues and at the sites of infection. The subunits of TCR heterodimers are covalently bonded and in the endoplasmic reticulum they associate with CD3 subunits to form functional TCR-CD3 complex. Lack of expression of any of the chains is sufficient to stop cell surface expression.

Product Info

Amount :	100 tests
Storage condition :	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human TCR alpha/beta (IP26) PE antibody (concentration in sample 1.67 µg/ml).




Figure 2: Flow cytometry multicolor surface staining pattern of human lymphocytes stained using anti-human CD3 (UCHT1) APC antibody (10 µl reagent / 100 µl of peripheral whole blood) and anti-human TCR alpha/beta (IP26) PE antibody (concentration in sample 1.67 µg/ml).




Figure 3: Separation of human CD3 positive TCR alpha/beta positive T cells (red-filled) from CD3 negative TCR alpha/beta negative lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human TCR alpha/beta (IP26) PE antibody (concentration in sample 1.67 $\mu\text{g/ml}$).