

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-1559: Monoclonal Antibody to Progesterone Receptor (Marker of Progestin Dependence)(Clone: PR501)

Clonality: Monoclonal PR501 **Clone Name:** Application: WB,IHC Reactivity: Human Gene: PGR Gene ID: 5241 **Uniprot ID:** P06401 Format: Purified **Alternative Name:** PGR.NR3C3

Isotype: Mouse IgG1, kappa

Immunogen Information: Recombinant human Progesterone Receptor protein

Description

This MAb is specific to progesterone receptor and shows minimal cross-reaction with other members of the family. Progesterone receptor is expressed as two major isoforms, PR-A (81kDa) and PR-B (116kDa). Expression of PgR has been suggested to reflect a intact estrogen regulatory machinery and therefore, predict better clinical response to endocrine therapy than ER alone.

Product Info

Amount: $100 \mu g$

Purification: Affinity Chromatography

Content: 100 μg in 500 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly


toxic.

Storage condition : Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Western Blot (1-2ug/ml for 2 hours at Room Temp); Immunohistochemistry (Formalin-fixed) (1-2ug/ml) (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);



Formalin-fixed, paraffin-embedded normal human Breast (left) and Invasive Ductal Carcinoma (right) stained with progesterone receptor Ab (PR501).