

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 42-1325: Anti-HSP40, YDJ1 Monoclonal Antibody (Clone: 1G10.H8)

Clonality: Monoclonal **Clone Name:** 1G10.H8 IP.ELISA.WB Application: Reactivity: Yeast

Conjugate: Unconjugated

Gene: YDJ1 Gene ID: 855661 **Uniprot ID:** P25491

**Alternative Name:** YDJ1,MAS5,YNL064C,N2418,YNL2418C

Isotype: Mouse IgG1 Kappa

Immunogen Information: Full length protein yeast HSP40 (YDJ1)

## **Description**

Human HSP40/DnaJ proteins comprise a large protein family, members of which feature the J domain (named after the bacterial Dnal protein). The I-domain spans the first 75 N-terminal amino acids and is separated from the C-terminal by a glycine/phenylalanine-rich domain. There are two main types of HSP40; type 1 DNAJ proteins including HDJ2 and yeast Ydjl; type II includes yeast Sis1 and human Hdj1. Whereas type I possesses a zinc finger domain which helps in the function of protein folding, type II does not. Members of the HSP40/DnaJ family play diverse roles in many cellular processes, such as folding, translocation, degradation and assembly of multi-protein complexes. HSP40 stimulates the ATPase activity of HSP70 which in turn causes conformational changes of the unfolded proteins. The HSP40-HSP70-unfolded protein complex further binds to co-chaperones Hip, Hop and HSP90 which leads to protein folding, or components of protein degradation machinery CHIP and BAG-1.

## **Product Info**

Amount: 100 µg

**Purification:** Protein G Purified

Content: 50% glycerol, 0.09% sodium azide

Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid Storage condition:

repeated freeze and thaw cycles.


## **Application Note**

WB (1:2000); optimal dilutions for assays should be determined by the user.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com



 $\label{thm:continuous} \mbox{Figure1: Mouse Anti-Hsp40 Antibody [1G10.H8] used in Western Blot (WB) on Yeast Cell lysates$