

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

42-1328: Anti-HSP40, YDJ1 Monoclonal Antibody (Clone: 1G10.H8) - ATTO 565(Discontinued)

Clonality: Monoclonal **Clone Name:** 1G10.H8 Application: WB.IP.ELISA Reactivity: Yeast Conjugate: **ATTO 565** Gene: YDJ1 Gene ID: 855661

P25491 **Alternative Name:** YDJ1,MAS5,YNL064C,N2418,YNL2418C

Isotype: Mouse IgG1 Kappa

Immunogen Information: Full length protein yeast HSP40 (YDJ1)

Description

Uniprot ID:

Human HSP40/DnaJ proteins comprise a large protein family, members of which feature the J domain (named after the bacterial DnaJ protein). The J-domain spans the first 75 N-terminal amino acids and is separated from the C-terminal by a glycine/phenylalanine-rich domain. There are two main types of HSP40; type 1 DNAJ proteins including HDJ2 and yeast YdjI; type II includes yeast Sis1 and human Hdj1. Whereas type I possesses a zinc finger domain which helps in the function of protein folding, type II does not. Members of the HSP40/DnaJ family play diverse roles in many cellular processes, such as folding, translocation, degradation and assembly of multi-protein complexes. HSP40 stimulates the ATPase activity of HSP70 which in turn causes conformational changes of the unfolded proteins. The HSP40-HSP70-unfolded protein complex further binds to co-chaperones Hip, Hop and HSP90 which leads to protein folding, or components of protein degradation machinery CHIP and BAG-1.

Product Info

Amount: 100 µg

Purification: Protein G Purified

Content: 50% glycerol, 0.09% sodium azide

Storage condition: Store the antibody at 4°C

Application Note

WB (1:2000); optimal dilutions for assays should be determined by the user.




Figure 1: Mouse Anti-Hsp40 Antibody [1G10.H8] used in Western Blot (WB) on Yeast Cell lysates