

## 42-1331: Anti-HSP40, YDJ1 Monoclonal Antibody (Clone : 1G10.H8) - ATTO 655(Discontinued)

|                                |  |
|--------------------------------|--|
| <b>Clonality :</b>             | Monoclonal                             |
| <b>Clone Name :</b>            | 1G10.H8                                |
| <b>Application :</b>           | WB,IP,ELISA                            |
| <b>Reactivity :</b>            | Yeast                                  |
| <b>Conjugate :</b>             | ATTO 655                               |
| <b>Gene :</b>                  | YDJ1                                   |
| <b>Gene ID :</b>               | 855661                                 |
| <b>Uniprot ID :</b>            | P25491                                 |
| <b>Alternative Name :</b>      | YDJ1,MAS5,YNL064C,N2418,YNL2418C       |
| <b>Isotype :</b>               | Mouse IgG1 Kappa                       |
| <b>Immunogen Information :</b> | Full length protein yeast HSP40 (YDJ1) |

### Description

Human HSP40/DnaJ proteins comprise a large protein family, members of which feature the J domain (named after the bacterial DnaJ protein). The J-domain spans the first 75 N-terminal amino acids and is separated from the C-terminal by a glycine/phenylalanine-rich domain. There are two main types of HSP40; type I DNAJ proteins including HDJ2 and yeast Ydj1; type II includes yeast Sis1 and human Hdj1. Whereas type I possesses a zinc finger domain which helps in the function of protein folding, type II does not. Members of the HSP40/DnaJ family play diverse roles in many cellular processes, such as folding, translocation, degradation and assembly of multi-protein complexes. HSP40 stimulates the ATPase activity of HSP70 which in turn causes conformational changes of the unfolded proteins. The HSP40-HSP70-unfolded protein complex further binds to co-chaperones Hip, Hop and HSP90 which leads to protein folding, or components of protein degradation machinery CHIP and BAG-1.

### Product Info

|                            |                                  |
|----------------------------|----------------------------------|
| <b>Amount :</b>            | 100 µg                           |
| <b>Purification :</b>      | Protein G Purified               |
| <b>Content :</b>           | 50% glycerol, 0.09% sodium azide |
| <b>Storage condition :</b> | Store the antibody at 4°C        |

### Application Note

WB (1:2000); optimal dilutions for assays should be determined by the user.




Figure1 : Mouse Anti-Hsp40 Antibody [1G10.H8] used in Western Blot (WB) on Yeast Cell lysates