

38-1047: Polyclonal antibody to Caspase-9

Clonality :	Polyclonal
Application :	FACS,WB
Reactivity :	Human
Gene :	CASP9
Gene ID :	842
Uniprot ID :	P55211
Format :	Purified
Alternative Name :	Apoptotic protease Mch-6, Apoptotic protease-activating factor 3, ICE-like apoptotic protease 6, MCH6
Isotype :	Rabbit IgG
Immunogen Information :	A synthetic peptide of Caspase-9 protein (amino acids 188-206 EKLRRRFSSLHFMVEVKGD) was used as the immunogen for this antibody

Description

Apoptosis, or programmed cell death, is a common property of all multicellular organisms. The current dogma of apoptosis suggests that the components of the core cell-death machinery are integral to cells and widely conserved across species. Caspases, a family of cysteinyl aspartate-specific proteases, are integral components of the cell death machinery (reviewed in Siegal, 2006; and Lavrik et al, 2005). They play a central role in the initiation and execution of apoptotic cell death and in inflammation. Caspases are typically divided into 3 major groups, depending on the structure of their prodomain and their function. Group I: inflammatory caspases (caspases 1, 4, 5, 11, 12, 14). Group II: initiator of apoptosis caspases (caspases 2, 8, 9). Group III: effector caspases (caspases 3, 6, 7). Caspases are synthesized as zymogens (inactive pro enzyme precursors which require a biochemical change to become active enzymes) with an N-terminal prodomain of variable length followed by a large subunit (p20) and a small subunit (p10). Caspases are activated through proteolytic cleavage at specific asparagine residues that are located within the prodomain, the p10, and p20 subunits. Activation results in the generation of mature active caspases that consist of the heterotetramer p20₂-p10₂. Active caspases mediate cell death and inflammation through cleavage of particular cellular substrates that are involved in these processes. The Caspase-9 polyclonal antisera recognizes the pro form of caspase-9 (approx. 50 kDa), and the large (approx. 35 kDa) and small (approx. 15 kDa) subunits of active/cleaved Caspase-9.

Product Info

Amount :	25 µg / 100 µg
Purification :	Protein A Chromatography
Content :	25 µg in 50 µl/100 µg in 200 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
Storage condition :	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Western blot analysis: 2-4 µg/ml, Flowcytometric analysis: 0.5-1/10⁶ cells




Figure-1: Western blot analysis of Caspase-9. Anti-Caspase-9 antibody (38-1047) was used at 1 $\mu\text{g/ml}$ on (1) PC3 and (2) HCT-116 lysates.




Figure-2: Intracellular staining of Anti-Caspase-9 in Jurkat cell line using 0.5 $\mu\text{g}/10^6$ cells. Green represent isotype control and red represent the Anti-caspase-9 antibody (38-1047). Goat Ant-Rabbit PE conjugated secondary was used as secondary antibody.