

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

42-1524: Anti-TCF7/TCF1 Polyclonal Antibody(Discontinued)

Clonality: Polyclonal
Application: IHC,WB
Reactivity: Human
Gene: TCF7
Gene ID: 6932
Uniprot ID: P36402
Format: Purified

Alternative Name: T-cell-specific transcription factor 1, Transcription factor 7, TCF7

Isotype: Rabbit IgG

Immunogen Information: Synthetic peptide from the C-terminal of Human TCF7 (284-384 aa) was used as immunogen.

Description

Transcriptional activator involved in T-cell lymphocyte differentiation. Necessary for the survival of CD4(+) CD8(+) immature thymocytes. Isoforms lacking the N-terminal CTNNB1 binding domain cannot fulfill this role. Binds to the T- lymphocyte-specific enhancer element (5'-WWCAAAG-3') found in the promoter of the CD3E gene. May also act as feedback transcriptional repressor of CTNNB1 and TCF7L2 target genes. TLE1, TLE2, TLE3 and TLE4 repress transactivation mediated by TCF7 and CTNNB1.

Product Info

Amount : 100 μ g **Purification :** Protein A

Content: PBS pH 7.4, 50% glycerol, 0.09% Sodium Azide

Storage condition : Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

WB (1:1000); IHC (1:50); optimal dilutions for assays should be determined by the user.




Fig-1: Western blot analysis of TCF-1 antibody. Anti- TCF-1 was used at 1:1000 dilution on Hela lysate.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com



Fig-2: Immunohistochemical analysis of TCF-1 in human Tonsil tissue using TCF-1 antibody at 1:50 dilutions.