w abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com

32-20139: Animal-Free Recombinant Human GDF-3(Discontinued)

Alternative Name : Growth/Differentiation Factor-3, Vgr-2, UNQ2222/PRO248

Description

Source:E.coliGDF-3 is a member of the TGF-Beta superfamily of growth and differentiation factors, and is highly homologous to GDF-9. Unlike most TGF-Beta family members, GDF-3 and GDF-9 are not disulfide-linked dimers. GDF-3 is expressed in adult bone marrow, spleen, thymus, and adipose tissue. The expression of GDF-3 is upregulated in high-fat-fed wild-type FABP4/aP2 null mice and was associated with obesity, but not with the related hyperglycemia/hyperinsulinemia that characterizes Type 2 diabetes. Recombinant Human GDF-3 is a 26.0 kDa non-disulfide-linked homodimer containing two 114 amino acid polypeptide chains.

Product Info

 Amount :
 5 μg / 20 μg

 Purification :
 Purity: >= 98% by SDS-PAGE gel and HPLC analyses.

 Content :
 This recombinant protein is supplied in lyophilized form.

 Amino Acid :
 AAIPVPKLSC KNLCHRHQLF INFRDLGWHK WIIAPKGFMA NYCHGECPFS LTISLNSSNY AFMQALMHAV DPEIPQAVCI PTKLSPISML YQDNNDNVIL RHYEDMVVDE CGCG

Application Note

Determined by its ability to inhibit induced alkaline phosphatase production by ATDC-5 chondrogenic cells. \tilde{A} \tilde{A} The \tilde{A} \tilde{A} ED₅₀ for this effect is 100-150 ng/ml.