∗ abeomics

36-1570: Monoclonal Antibody to ACTH (Adrenocorticotrophic Hormone) (Pituitary Marker)(Clone : SPM333)

| Clonality : | Monoclonal |
|-----------------------|---|
| Clone Name : | SPM333 |
| Application : | ELISA,FACS,IF,IHC |
| Reactivity : | Human, Mouse, Rat |
| Gene : | РОМС |
| Gene ID : | 5443 |
| Uniprot ID : | P01189 |
| Format : | Purified |
| Alternative Name : | РОМС |
| Isotype : | Mouse IgG1, kappa |
| Immunogen Information | Synthetic peptide corresponding to aa1-24 of human ACTH |

Description

ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by the anterior pituitary. This MAb is specific to Synacthen (aa1-24 of ACTH); does not react with CLIP (aa17-39 of ACTH). POMC (pro-opiomelanocortin or corticotropinlipotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific posttranslational processing by convertases. POMC is cleaved into ten hormone chains named NPP, ACTH, alpha-MSH (Melanocyte Stimulating Hormone), beta-MSH, gamma-MSH, CLIP (corticotropin-like intermediary peptide), Lipotropin-beta, Lipotropin-gamma, betaendorphin and Met-enkephalin. ACTH is also produced by cells of immune system (T-cells, B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pituitary tumors and the study of pituitary disease. It reacts with ACTH-producing cells (corticotrophs). It also may react with other tumors (e.g. some small cell carcinomas of the lung) causing paraneoplastic syndromes by secreting ACTH.

Product Info

| Amount : | 100 µg |
|---------------------|---|
| Purification : | Affinity Chromatography |
| Content : | 100 μg in 500 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic. |
| Storage condition : | Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles. |

Application Note

ELISA (For coating, order Ab without BSA); Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT) (No special pretreatment is required for staining of formalin-fixed tissues)