

## 32-20316: Recombinant Human NOV(Discontinued)

**Reactivity :** Chicken, Human, Mouse

### Description

**Source:**E.coli

NOV is a member of the CCN family of secreted, cysteine-rich regulatory proteins. The full-length NOV protein contains four structural domains that confer distinct, and sometimes opposing, biological activities. Elevated expression of NOV is associated with certain tumors, including Wilm's tumor and most neuroblastomas. However, in other tumor types and certain cancer cell lines, increased tumorigenicity and proliferation is correlated with decreased NOV expression. Additionally, NOV induces cell adhesion and cell migration by signaling through specific cell-surface integrins, and by binding to heparin sulfate proteoglycans and to fibulin 1C. NOV has also been reported to exert proangiogenic activities. Recombinant Human NOV is a 36.2 kDa protein containing 331 amino acid residues. It is composed of four distinct structural domains (modules): the IGF binding protein (IGFBP) domain; the von Willebrand Factor C (VWFC) domain; the Thrombospondin type-1 (TSP type-1) domain; and a C-terminal cysteine knot-like domain (CTCK).

### Product Info

**Amount :** 5 µg / 20 µg

**Purification :** Purity: >= 95% by SDS-PAGE gel and HPLC analyses.

**Content :** This recombinant protein is supplied in lyophilized form.

**Amino Acid :** MQVAATQRCP PQCPGRCPAT PPTCAPGVRA VLDGCSCCLV CARQGESCS DLEPCDESSG  
LYCDRSADPS NQTGICTAVE GDNCVFDGVI YRSGEKFQPS CKFQCTCRDG QIGCVPRCQL DVLLPEPNCP  
APRKVEVPGE CCEKWICGPD EEDSLGGLTL AAYRPEATLG VEVSDSSVNC IEQTTEWTAC SKSCGMGFST  
RVTNRRNQCE MLKQTRLCMV RPCEQEPEQP TDKKGGKCLR TKKSLKAIHL QFKNCTSLHT YKPRFCGVCS  
DGRCTPHNT KTIQAEFQCS PGQIVKKPVM VIGTCTHTN CPKNNEAFLQ EELKTTTRGK M

### Application Note

Determined by a cell proliferation assay using BALB/c 3T3 cells. The expected ED<sub>50</sub> for this effect is 1.0-2.0 µg/ml