

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com

32-20383: Animal-Free Recombinant Human IL-17E(Discontinued)

Alternative Name : Interleukin-17E, IL-25

Description

Source:E.coli

IL-17E is a disulfide-linked homodimer of two 145 amino acid polypeptide chains. It belongs to the IL-17 family of structurally-related cytokines that share a highly conserved C-terminal region, but differ from one another in their N-terminal regions and in their distinct biological roles. The six known members of this family, IL-17A through IL-17F, are secreted as homodimers. IL-17E stimulates secretion of IL-8, and induces activation of the transcription factor NF-kB in cells that express the IL-17BR receptor. Recombinant Human IL-17E is a 33.8 kDa disulfide-linked homodimer of two 146 amino acid polypeptide chains.

Product Info

 Amount :
 5 μg / 25 μg

 Purification :
 Purity:>= 98% by SDS-PAGE gel and HPLC analyses.

 Content :
 This recombinant protein is supplied in lyophilized form.

 Amino Acid :
 MYSHWPSCCP SKGQDTSEEL LRWSTVPVPP LEPARPNRHP ESCRASEDGP LNSRAISPWR YELDRDLNRL PQDLYHARCL CPHCVSLQTG SHMDPRGNSE LLYHNQTVFY RRPCHGEKGT HKGYCLERRL YRVSLACVCV RPRVMG

Application Note

Determined by its ability to induce IL-8 in human PBMCs using a concentration range of 10-100 ng.