

32-20407: Recombinant Human PDGF-BB(Discontinued)

Reactivity : Cow, Guinea Pig, Human, Monkey, Mouse, Pig, Rat

Alternative Name : Platelet-Derived Growth Factor-BB, Glioma-derived growth factor (GDGF), Osteosarcoma-derived Growth Factor (ODGF)

Description

Source: *E.coli* PDGFs are disulfide-linked dimers consisting of two 12.0-13.5 kDa polypeptide chains, designated PDGF-A and PDGF-B chains. The three naturally occurring PDGFs, PDGF-AA, PDGF-BB and PDGF-AB, are potent mitogens for a variety of cell types, including smooth muscle cells, connective tissue cells, bone and cartilage cells, and some blood cells. The PDGFs are stored in platelet Alpha -granules, and are released upon platelet activation. The PDGFs are involved in a number of biological processes, including hyperplasia, chemotaxis, embryonic neuron development, and respiratory tubule epithelial cell development. Two distinct signaling receptors used by PDGFs have been identified and named PDGFR-Alpha and PDGFR-Beta . PDGFR-Alpha is a high-affinity receptor for each of the three PDGF forms. On the other hand, PDGFR-Beta interacts with only PDGF-BB and PDGF-AB. Recombinant Human PDGF-BB is a 24.3 kDa disulfide-linked homodimer of two β chains (218 total amino acids). .

Product Info

Amount : 2 μ g / 10 μ g

Purification : Purity: \geq 98% by SDS-PAGE gel and HPLC analyses.

Content : This recombinant protein is supplied in lyophilized form.

Amino Acid : SLGSLTIAEP AMIAECKTRT EVFEISRRLI DRTNANFLVW PPCVEVQRCS GCCNNRNVQC RPTQVQLRPV
QVRKIEIVRK KPIFKKATVT LEDHLACKCE TVAAARPVT

Application Note

Determined by the dose-dependent stimulation of the proliferation of Balb/c 3T3 cells. The expected ED_{50} for this effect is 1.0-3.0 ng/ml.