

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-20476: Recombinant Human M-CSF(Discontinued)

**Reactivity:** Human, Monkey, Mouse, Rat

Alternative Name: Macrophage Colony Stimulating Factor, CSF-1, MGI-IM

## **Description**

**Source:E.coli**M-CSF is a potent hematopoietic factor produced by a variety of cells, including lymphocytes, monocytes, fibroblasts, endothelial cells, myoblasts and osteoblasts. It is a key regulator of cellular proliferation, differentiation, and survival for blood monocytes, tissue macrophages, and their respective progenitor cells. M-CSF has been shown to play important roles in modulating dermal thickness and fertility. M-CSF is clinically used in the treatment of infection, malignancies and atherosclerosis. It facilitates hematopoietic recovery after bone marrow transplantation. Human M-CSF is reactive in murine systems, but the murine molecule exhibits no activity on human cells. Recombinant Human M-CSF is a 36.8 kDa homodimeric protein consisting of two 159 amino acid polypeptide subunits.

## **Product Info**

Amount:  $2 \mu g / 10 \mu g$ 

**Purification:** Purity:>= 98% by SDS-PAGE gel and HPLC analyses. **Content:** This recombinant protein is supplied in lyophilized form.

Amino Acid: MEEVSEYCSH MIGSGHLQSL QRLIDSQMET SCQITFEFVD QEQLKDPVCY LKKAFLLVQD IMEDTMRFRD

NTPNAIAIVQ LQELSLRLKS CFTKDYEEHD KACVRTFYET PLQLLEKVKN VFNETKNLLD KDWNIFSKNC

NNSFAECSSQ GHERQSEGS

## **Application Note**

The $\tilde{A}$  $\|\hat{A}$  ED<sub>50</sub> $\tilde{A}$  $\|\hat{A}$  was determined by the dose-dependent stimulation of the proliferation of murine M-NSF-60 cells <=1 ng/ml, corresponding to a specific activity of>= 1 x 10<sup>6</sup> $\tilde{A}$  $\|\hat{A}$  units/mg. $\tilde{A}$  $\|\hat{A}$