

32-20491: Recombinant Human sRANK Receptor(Discontinued)

Reactivity : Human, Mouse

Alternative Name : TNFRSF11A, ODFR (osteoclast differentiation factor receptor), ODAR (osteoclast differentiation and activation receptor), TRANCE Receptor

Description

Source: E.coli RANKL and RANK are members of the TNF superfamily of ligands and receptors that play an important role in the regulation of specific immunity and bone turnover. RANK (receptor) was originally identified as a dendritic cell-membrane protein, which, by interacting with RANKL, augments the ability of dendritic. These dendritic cells then stimulate naïve T-cell proliferation, and promote the survival of RANK + T-cells. RANK is also expressed in a variety of tissues, including skeletal muscle, thymus, liver, colon, small intestine, and adrenal gland. The RANK/RANKL interaction is important in the regulation of osteoclastogenesis, and in dendritic cell-mediated T-cell immune responses. Impairments in RANK signaling have been implicated in the induction of expansile osteolysis and Paget's disease of bone (PDB2). Recombinant Human sRANK Receptor is a 19.3 kDa polypeptide containing the TNFR-homologous, cysteine-rich portion of the extracellular domain of RANK receptor (175 amino acid residues).

Product Info

Amount : 20 µg / 100 µg

Purification : Purity: >= 98% by SDS-PAGE gel and HPLC analyses.

Content : This recombinant protein is supplied in lyophilized form.

Amino Acid : MQIAPPCTSE KHYEHLGRCC NKCEPGKYMS SKCTTTSDSV CLPCGPDEYL DSWNEEDKCL LHKVCDTGKA
LVAVVAGNST TPRRCAC TAG YHWSQDCECC RRNTECAPGL GAQHPLQLNK DTVCKPCLAG
YFSDAFSSTD KCRPWTNCTF LGKRVEHHGT EKSDAVCSSS LPARK

Application Note

Determined by its ability to inhibit sRANKL induced NF-κB in RAW264.7 cells in the absence of any cross-linking. The expected ED_{50} for this effect in the presence of 15 ng/ml of recombinant sRANKL, is 30-50 ng/ml.