

## 36-1580: Monoclonal Antibody to DOG-1 / TMEM16A / ANO1 (Gastrointestinal Stromal Tumor Marker)(Clone : SPM580)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	SPM580
<b>Application :</b>	IHC
<b>Reactivity :</b>	Human
<b>Gene :</b>	ANO1
<b>Gene ID :</b>	55107
<b>Uniprot ID :</b>	Q5XXA6
<b>Format :</b>	Purified
<b>Alternative Name :</b>	ANO1,DOG1,ORAOV2,TAOS2,TMEM16A
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Recombinant human DOG-1 protein

### Description


Expression of DOG-1 protein is elevated in the gastrointestinal stromal tumors (GISTs), c-kit signaling-driven mesenchymal tumors of the GI tract. DOG-1 is rarely expressed in other soft tissue tumors, which, due to appearance, may be difficult to diagnose. Immunoreactivity for DOG-1 has been reported in 97.8 percent of scorable GISTs, including all c-kit negative GISTs. Overexpression of DOG-1 has been suggested to aid in the identification of GISTs, including Platelet-Derived Growth Factor Receptor Alpha mutants that fail to express c-kit antigen. The overall sensitivity of DOG1 and c-kit in GISTs is nearly identical: 94.4% vs. 94.7%.

### Product Info

<b>Amount :</b>	100 µg
<b>Purification :</b>	Affinity Chromatography
<b>Content :</b>	100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
<b>Storage condition :</b>	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);



Formalin-fixed, paraffin-embedded human GIST stained with DOG1 Monoclonal Antibody (SPM580).