

## 32-20503: Recombinant Murine TRAIL(Discontinued)

**Reactivity :** Human, Mouse

**Alternative Name :** TNF-Related Apoptosis-Inducing Ligand, TNFSF10, TL2

### Description

**Source:** **E.coli**TRAIL is a cytotoxic protein, which activates rapid apoptosis in tumor cells, but not in normal cells. TRAIL-induced apoptosis is achieved through binding to two death-signaling receptors, DR4 and DR5. These receptors belong to the TNFR superfamily of transmembrane proteins, and contain a cytoplasmic "death domain", which activates the cell's apoptotic machinery. Recombinant Murine TRAIL is a 174 amino acid polypeptide (20.0 kDa), consisting of the TNF-homologous portion of the extracellular domain of the full length TRAIL protein.

### Product Info

**Amount :** 10 µg / 50 µg

**Purification :** Purity: >= 95% by SDS-PAGE gel and HPLC analyses.

**Content :** This recombinant protein is supplied in lyophilized form.

**Amino Acid :** MRGGRPQKVA AHITGITRRS NSALIPISKD GKTLGQKIES WESSRKGHSF LNHVLFNRNGE LVIEQEGLYY  
IYSQTYFRFQ EAEDASKMVS KDKVRTKQLV QYIYKYTSYP DPIVLMKSAR NSCWSRDAEY GLYSIQGGGL  
FELKKNDRIF VSVTNEHLMD LDQEASFFGA FLIN

### Application Note

Assay#1: Determined by the dose-dependent stimulation of MIP-2 production by mouse spleen cells using a concentration range of 10-100 ng/ml. Assay#2: Measured by its ability to induce apoptosis in LN-18 cells (human glioblastoma cells). The expected ED<sub>50</sub> for this effect is 40.0-60.0 ng/ml.