

32-20600: Recombinant Human PECAM-1(Discontinued)

Alternative Name : Platelet endothelial cell adhesion molecule, CD31 antigen, EndoCAM

Description

Source:HEK293 cells

PECAM is transmembrane glycoprotein that belongs to the Ig-related superfamily of adhesion molecules. It is highly expressed at endothelial cell junctions, and is also expressed in platelets and most leukocyte sub-types. The primary function of PECAM-1 is the mediation of leukocyte-endothelial cell adhesion and signal transduction. PECAM-1 has been implicated in the pathogenesis of various inflammation-related disorders, including thrombosis, multiple sclerosis (MS), and rheumatoid arthritis. The human PECAM-1 gene codes for a 738 amino acid transmembrane glycoprotein that contains a 118 amino acid cytoplasmic domain, a 19 amino acid transmembrane domain, and a 574 amino acid extracellular domain. Recombinant Human PECAM-1 is a 572 amino acid glycoprotein comprising the extracellular domain of PECAM-1. Monomeric glycosylated PECAM-1 migrates at an apparent molecular weight of approximately 80.0-95.0 kDa by SDS-PAGE analysis under reducing conditions. The calculated molecular weight of Recombinan Human PECAM-1 is 64.3 kDa.

Product Info

Amount : 10 µg / 50 µg

Purification : Purity:>= 97% by SDS-PAGE gel and HPLC analyses.

Content : This recombinant protein is supplied in lyophilized form.

Amino Acid : ENSFTINSV DMKSLPDWTV QNGKNLTLQC FADVSTTSHV KPQHQLFYK DDVLFYNISS MKSTESYFIP
EVRIYDSGT Y KCTVIVNNKE KTTAEYQLLV EGVPSRVTL DKKEAIQGGI VRVNCSPVEE KAPIHFTIEK
LELNEKMKV KL KREKNSRDQN FVILEFPVEE QDRVLSFRCQ ARIISGIHQ TSESTKSELV TVTESFSTPK
FHISPTGMIM EGAQLHIKCT IQVTHLAQEF PEIIIQKDKA IVAHNRHGK AVYSVMAMVE HSGNYTCKVE
SSRISKVSSI VVNITELFSK PELESSFTHL DQGERLNLSC SIPGAPPANF TIQKEDTIVS QTQDFTKIAS
KSDSGTYICT AGIDKVVKKS NTVQIVVCEM LSQPRISYDA QFEVIKQTI EVRCESISGT LPISYQLLKT
SKVLENSTKN SNDPAVFKDN PTEDVEYQCV ADNCHSHAKM LSEVLRVKVI APVDEVQISI LSSKVVESGE
DIVLQCAVNE GSGPITYKFY REKEGKPFYQ MTSNATQAFW TKQKASKEQE GEYCYTAFNR ANHASSVPRS
KILTRVILA PWK

Application Note

Determined by its ability to support the adhesion of activated Jurkat cells. The expected ED_{50} for this effect is 1.0 -1.5 µg/ml.