

32-20619: Recombinant Human IL-12 p70 (HEK293 derived)(Discontinued)

Alternative Name :

Interleukin-12, NKSF, CTL Maturation Factor (TCMF), Cytotoxic Lymphocyte Maturation Factor (CLMF), TSF p35: Interleukin-12 subunit alpha, IL-12 subunit p35, IL-12A, Cytotoxic Lymphocyte Maturation Factor 35 kDa subunit (CLMF p35), NK cell Stimulating Factor Chain 1 p40: Interleukin-12 subunit beta, IL-12 subunit p40, IL-12B, Cytotoxic Lymphocyte Maturation Factor 40 kDa subunit (CLMF p40), NK cell Stimulating Factor Chain 2

Description

Source:HEK293 cells

IL-12 is a potent regulator of cell-mediated immune responses and it induces IFN-Gamma production by NK and T cells. It is produced by activated monocytes/macrophage cells, B lymphocytes and connective tissue-type mast cells. Among its biological activities, IL-12 promotes the growth and activity of activated NK, CD4+ and CD8+ cells, and induces the development of IFN-Gamma-producing Th1 cells. The Recombinant Human IL-12 p70, derived from HEK293 cells, is a heterodimeric glycoprotein consisting of disulfide-linked p35 and p40 subunits (503 total amino acid residues). The calculated molecular weight of Human IL-12 p70 is 57.2 kDa; however, due to glycosylation, it migrates at an apparent molecular weight of 65-70 kDa based on SDS-PAGE gel, under non-reducing conditions.

Product Info

Amount : 2 µg / 10 µg

Purification : Purity: >= 98% by SDS-PAGE gel and HPLC analyses.

Content : This recombinant protein is supplied in lyophilized form.

Amino Acid :
p35 Subunit: RNLPVATPDP GMFPC LHHSQ NLLRAVSNML QKARQTLEFY PCTSEEIDHE DITKDKTSTV
EACLPLELTK NESCLNSRET SFITNGSCLA SRKTSFMMAL CLSSYEDLK MYQVEFKTMN AKLLMDPKRQ
IFLDQNMLAV IDELMQALNF NSETVPQKSS LEEPDFYKTK IKLCILLHAF RIRAVTIDRV MSYLNAS p40
Subunit: IWELKKDVYV VELD WYPDAP GEMVVLTCDT PEEDGITWTL DQSEVLGSG KTLTIQVKEF
GDAGQYTCHK GGEVLSHLL LLHKEDGIW STDILKDQKE PKNKTFLRCE AKNYSGRFTC WWLTTISTDL
TFSVKSSRGS SDPQGVTCGA ATLSAERVRG DNKEYEYSVE CQEDSACPAE EESLPIEMV DAVHKLKYEN
YTSSFFIRDI IKPDPPKNLQ LKPLKNSRQV EVSWEYPDTW STPHSYFSLT FCVQVQGKSK REKKDRVFTD
KTSATVICRK NASISVRAQD RYSSSWSEW ASVPCS

Application Note

Determined by its ability to increase IFN-Gamma production by anti-TCR mAb-stimulated PBMCs.