

14-516ACL: VISTA Stable Cell Line

Application : Functional Assay

Description

VISTA Stable Cell Line is a stably transfected CHO-K1 cell line which expresses human V-type immunoglobulin domain-containing suppressor of T-cell activation (VISTA, also known as VSIR, B7-H5, PD-1H, and SISP1).

Sequence data: hVISTA (accession number NP_071436)

```
MGVPTALEAGSWRWGSLLFALFLAASLGPVAAFVATPYSLYVC  
PEGQNVTLTCRLLGPVDKGHDVTFYKTWYRSSRGEVQTCSERRPIRNLTFQDLHLHHG  
GHQAANTSHDLAQRHGLESADHHGNFSITMRNLTLDSGLYCCLVVEIRHHHSEHRV  
HGAMELQVQTGKDAPSNVYVYSSSQDSENITAAALATGACIVGILCLPLILLVYKQ  
RQAASNRRAQELVRMDSNIQGIENPGFEASPPAQGIPEAKVRHPLSYVAQRQPSESGR  
HLLSEPSTPLSPPGPGDVFFPSLDPVPDSPNFEVI
```

Product Info

Amount : 1 Vial
Content : Each vial contains $2 \sim 3 \times 10^6$ cells in 1 ml of 90% FBS + 10% DMSO
Storage condition : Immediately upon receipt, store in liquid nitrogen.

Application Note

Application:

- Screen for antibodies of human VISTA through Flow Cytometry.

Culture conditions:

Cells should be grown at 37°C with 5% CO₂ using DMEM medium (w/ L-Glutamine, 4.5g/L Glucose and Sodium Pyruvate) supplemented with 10% heat-inactivated FBS and 1% Pen/Strep, plus 10 µg/ml of Blasticidin.

It is recommended to quickly thaw the frozen cells upon receipt or from liquid nitrogen in a 37°C water-bath, transfer to a tube containing 10 ml of growth medium without Blasticidin, spin down cells, resuspend cells in pre-warmed growth medium without Blasticidin, transfer resuspended cells to T25 flask and culture in 37°C-CO₂ incubator.

Leave the T25 flask in the incubator for 1~2 days without disturbing or changing the medium until cells completely recover viability and become adherent. Once cells are over 90% adherent, remove growth medium and passage the cells through trypsinization and centrifugation. At first passage, switch to growth medium containing Blasticidin. Cells should be split before they reach complete confluence.

To passage the cells, detach cells from culture vessel with Trypsin/EDTA, add complete growth medium and transfer to a tube, spin down cells, resuspend cells and seed appropriate aliquots of cells suspension into new culture vessels. Subcultivation ration = 1:10 to 1:20 weekly. To achieve satisfactory results, cells should not be passaged over 16 times.

LIMITED USE RESTRICTIONS:

THIS PRODUCT IS SOLELY FOR IN VITRO RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

By use of this product, user agrees to be bound by the terms of this limited use statement.

This product is solely for Internal Research Purposes and not for Commercial Purposes. Commercial Purposes include, but are not limited to (1) use of the cell line in manufacturing; (2) use of the cell line to provide a service, information or data; (3) use of the cell line for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the cell line whether or not such cell lines are resold for use in research. The buyer cannot sell, give or otherwise transfer this product to a third party.

Commercial License Agreement is available for non-research use if applicable. Please contact Abeomics (info@abeomics.com).




Fig-1: Detection of human VISTA in the CHO-K1/VISTA stable cell line by Flow Cytometry [Cell surface staining]. CHO-K1 cells (Green); CHO-K1/VISTA cells (Red).