

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 44-1125: Anti-TTF-1 Monoclonal Antibody (Clone:IHC414)(Discontinued)

Clonality: Monoclonal **Clone Name:** IHC414 Application: IHC Reactivity: Human Gene: NKX2-1 Gene ID: 7080 **Uniprot ID:** P43699 Format: **Purified** 

Alternative Name: NKX2-1, NKX2A, TITF1, TTF1, Homeobox protein NK-2 homolog A, Thyroid nuclear factor 1, Thyroid

transcription factor 1

## **Description**

Thyroid Transcription Factor 1 (TTF-1) is present in diencephalon, lung, and thyroid. Anti-TTF-1 stains thyroid and thyroid-derived tumors, and is therefore used for distinguishing lung adenocarcinoma from germ cell tumors, malignant mesothelioma, and metastatic carcinomas from organs other than the thyroid. It is also useful for distinguishing small cell lung carcinoma from lymphoid infiltrates, and pulmonary from non-pulmonary adenocarcinomas in malignant effusions. The ability to distinguish between pulmonary and non-pulmonary adenocarcinomas is particularly useful in identifying tumors that have metastasized to the brain.

## **Product Info**

**Amount:** 0.1 ml / 1 ml

**Purification :** Protein A/G Chromatography

**Storage condition :** Store at 2°C - 8°C.

## **Application Note**

Recommended dilutions: Immunohistochemical analysis: 1:100 - 1:200. However, this need to be optimized based on the research applications.




Figure 1: Immunohistochemical analysis of TTF-1 (IHC414) on Thyroid Gland