

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

37-1038: Human bFGF / FGF2 Recombinant Protein(Discontinued)

Reactivity: Human

Alternative Name: BFGF Protein, FGF-2 Protein, FGFB Protein, HBGF-2 Protein,

Description

Source : E. coli

Basic fibroblast growth factor (bFGF), also known as FGF2, is a member of the fibroblast growth factor (FGF) family. It is a highly specific chemotactic and mitogenic factor for many cell types, appears to be involved in remodeling damaged tissue, such as ulcer healing, vascular repair, traumatic brain injury (TBI). bFGF is a critical component of human embryonic stem cell culture medium. In addition, bFGF protein is a heparin-binding cationic protein involved in a variety of pathological conditions including angiogenesis and solid tumour growth. Thus, bFGF is regarded as a target for cancers chemopreventive and therapeutic strategies. bFGF/FGF2 Protein & Antibody Products

Product Info

Amount: Human bFGF / FGF2 Recombinant Protein(Discontinued) / 50 μg

Purification: > 95 % as determined by SDS-PAGE

Formulation Lyophilized from sterile 20mM Tris, pH 7.4

Content: Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before

lyophilization.

Storage condition:

Storage condition:

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be

aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Amino Acid: Pro143-Ser288

Application Note

Measured in a cell proliferation assay using BALB/c 3T3 mouse embryonic fibroblasts. The ED50 for this effect is typically 0.02-0.1 ng/ml

Endotoxin :< 1.0 EU per μ g protein as determined by the LAL method.

Other pack size also available.




Fig 1: Human bFGF / FGF2 Recombinant Protein



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig 2: Human bFGF / FGF2 Recombinant Protein measured in a cell proliferation assay using BALB/c 3T3 mouse embryonic fibroblasts.