

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

### 37-1090: Human CD147 / EMMPRIN / Basigin Recombinant Protein (His Tag)(Discontinued)

**Reactivity:** Human

Alternative Name: 5F7 Protein, Basigin Protein, CD147 Protein, EMMPRIN Protein, M6 Protein, OK Protein, TCSF Protein,

## **Description**

#### Source: HEK293 Cells

CD147/EMMPRIN (Extracellular Matrix Metalloproteinase Inducer), also known as Basigin (BSG), is a transmembrane glycoprotein with different forms resulted from different modes of glycosylation and N-terminal sequence variants. It is a member of the immunoglobulin superfamily with homology to both the immunoglobulin V domain and MHC class II antigen beta-chain. This protein play important roles in variety of events including spermatogenesis, embryo implantation, neural network formation. CD147 induces the production and release of matrix metalloproteinases (MMP) in the surrounding mesenchymal cells and tumor cells, and thereby promotes invasion, metastasis, growth and survival of malignant cells. Furthermore, CD147 also serves as a receptor for extracellular cyclophilinthe and its association with integrins might be important in signal transduction. Recently, CD147 displays increased expression in many cancers, and it has been previously demonstrated to participate in cancer metastasis and progression. Thus, CD147 and its antibody are used as an effective treatment for malignant cancers.

#### **Product Info**

Amount: Human CD147 / EMMPRIN / Basigin Recombinant Protein (His Tag)(Discontinued) / 100 µg

**Purification:** > 97 % as determined by SDS-PAGE

Formulation Lyophilized from sterile PBS, pH 7.4

**Content :** Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before

lyophilization.

**Storage condition :** Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be

aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Amino Acid: Met1-His205

# **Application Note**

Endotoxin :< 1.0 EU per µg of the protein as determined by the LAL method




Fig 1: Human CD147 / EMMPRIN / Basigin Recombinant Protein (His Tag)