

37-1098: Human Erythropoietin / EPO Recombinant Protein (Fc Tag)(Discontinued)

Reactivity : Human

Alternative Name : EP Protein, Erythropoietin Protein, MVCD2 Protein,

Description

Source : HEK293 Cells

Erythropoietin is a member of the EPO / TPO family. It is a secreted, glycosylated cytokine composed of four alpha helical bundles. Erythropoietin can be found in the plasma and regulates red cell production by promoting erythroid differentiation and initiating hemoglobin synthesis. It also has neuroprotective activity against a variety of potential brain injuries and antiapoptotic functions in several tissue types. Erythropoietin is the principal hormone involved in the regulation of erythrocyte differentiation and the maintenance of a physiological level of circulating erythrocyte mass. It is produced by kidney or liver of adult mammals and by liver of fetal or neonatal mammals. Genetic variation in erythropoietin is associated with susceptibility to microvascular complications of diabetes type 2. These are pathological conditions that develop in numerous tissues and organs as a consequence of diabetes mellitus. They include diabetic retinopathy, diabetic nephropathy leading to end-stage renal disease, and diabetic neuropathy. Diabetic retinopathy remains the major cause of new-onset blindness among diabetic adults. It is characterized by vascular permeability and increased tissue ischemia and angiogenesis. It has a longer circulating half-life in vivo. Erythropoietin is being much misused as a performance-enhancing drug in endurance athletes.

Product Info

Amount : Human Erythropoietin / EPO Recombinant Protein (Fc Tag)(Discontinued) / 20 µg

Purification : > 95 % as determined by SDS-PAGE.

Content : Formulation Lyophilized from sterile PBS, pH 7.4.
Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.

Storage condition : Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Amino Acid : Met1-Arg193

Application Note

Endotoxin :< 1.0 EU per μg protein as determined by the LAL method.

Other pack size also available.




Fig 1: Human Erythropoietin / EPO Recombinant Protein (Fc Tag)