

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

37-1164: Mouse ALK-3 / BMPR1A Recombinant Protein (His Tag)(Discontinued)

Reactivity: Mouse

Alternative Name: 1110037I22Rik Protein, Mouse; ALK3 Protein, Mouse; AU045487 Protein, Mouse; Bmpr Protein, Mouse;

BMPR-IA Protein, Mouse

Description

Source: HEK293 Cells

Activin receptor-Like Kinase 3 (ALK-3), also known as Bone Morphogenetic Protein Receptor, type IA (BMPR1A), is a type I receptor for bone morphogenetic proteins (BMPs) which belong to the transforming growth factor beta (TGF-beta) superfamily. The BMP receptors form a subfamily of transmembrane serine/threonine kinases including the type I receptors BMPR1A and BMPR1B and the type II receptor BMPR2. ALK-3/BMPR1A is expressed in the epithelium during branching morphogenesis. Deletion of BMPR1A in the epithelium with an Sftpc-cre transgene leads to dramatic defects in lung development. ALK-3 and ALK-6 share a high degree of homology, yet possess distinct signaling roles. The transforming growth factor (TGF)-beta type III receptor (TbetaRIII) enhanced both ALK-3 and ALK-6 signaling. TbetaRIII associated with ALK-3 primarily through their extracellular domains, whereas its interaction with ALK-6 required both the extracellular and cytoplasmic domains. ALK-3 plays an essential role in the formation of embryonic ventral abdominal wall, and abrogation of BMP signaling activity due to gene mutations in its signaling components could be one of the underlying causes of omphalocele at birth. The type IA BMP receptor, ALK-3 was specifically required at mid-gestation for normal development of the trabeculae, compact myocardium, interventricular septum, and endocardial cushion. Cardiac muscle lacking ALK-3 was specifically deficient in expressing TGFbeta2, an established paracrine mediator of cushion morphogenesis. Hence, ALK-3 is essential, beyond just the egg cylinder stage, for myocyte-dependent functions and signals in cardiac organogenesis.

Product Info

Amount: 3 / BMPR1A Recombinant Protein (His Tag)(Discontinued) / 200 µg

Purification: > 97 % as determined by SDS-PAGE

Formulation Lyophilized from sterile PBS, pH 7.4

Content: Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before

lyophilization.

Storage condition:

Storage condition:

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be

aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Amino Acid: Met1-Arg152

Application Note

Measured by its ability to inhibit recombinant human BMP4 induced activity in MC3T3-E1 Mouse osteoblastic cells. The ED50 for this effect is typically 0.5-2 $\tilde{A} \square \hat{A} \mu g/ml$ in the presence of 50 ng/ml of recombinant human BMP4. Endotoxin :< 1.0 EU per $\tilde{A} \square \hat{A} \mu g$ of the protein as determined by the LAL method




Fig 1: Mouse ALK-3 / BMPR1A Recombinant Protein (His Tag)



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig 2: Mouse ALK-3 / BMPR1A Recombinant Protein (His Tag) measured by its ability to inhibit recombinant human BMP4 induced activity in MC3T3-E1 Mouse osteoblastic cells.