

## 37-1171: Mouse CD38 Recombinant Protein (His Tag)(Discontinued)

**Reactivity :** Mouse

**Alternative Name :** ADPRC 1-rs1 Protein, Mouse; ADPRC1 Protein, Mouse; Cd38 Protein, Mouse; Cd38-rs1 Protein, Mouse; I-19 Protein, Mouse

### Description

#### Source : HEK293 Cells

The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 32 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. Cluster of differentiation 38 (CD38), also known as ADP-ribosyl cyclase, is a glycoprotein found on the surface of many immune cells (white blood cells), including CD4+, CD8+, B and natural killer cells. It shares several characteristics with ADP-ribosyl cyclase 2 CD157. CD38 is a multifunctional ectoenzyme that catalyzes the synthesis and hydrolysis of cyclic ADP-ribose (cADPR) from NAD<sup>+</sup> to ADP-ribose. It also functions in cell adhesion, signal transduction and calcium signaling. CD38 has been used as a prognostic marker in leukemia. It can also be used to identify plasma cells.

### Product Info

**Amount :** Mouse CD38 Recombinant Protein (His Tag)(Discontinued) / 20 µg

**Purification :** > 97 % as determined by SDS-PAGE

**Content :** Formulation Lyophilized from sterile 20mM MES, 0.15M NaCl, pH 6.5  
Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.

**Storage condition :** Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

**Amino Acid :** Leu45-Thr304

### Application Note

Measured by its ability to convert the substrate nicotinamide guanine dinucleotide (NGD+) to cyclic GDPribose. The specific activity is >50,000pmols/min/ug.

Endotoxin :< 1.0 EU per µg of the protein as determined by the LAL method.

Other pack size also available.




Fig 1: Mouse CD38 Recombinant Protein (His Tag)