

## 37-1174: Mouse ICOS / AILIM / CD278 Recombinant Protein (Fc Tag)(Discontinued)

**Reactivity :** Mouse  
**Alternative Name :** AILIM Protein, Mouse; CCLP Protein, Mouse; CRP-1 Protein, Mouse; H4 Protein, Mouse; Ly115 Protein, Mouse

### Description

#### Source : HEK293 Cells

Inducible costimulator (ICOS), also called AILIM (activation-inducible lymphocyte immunomediatory molecule) is a cell-surface receptor, and belongs to the CD28 family of immune costimulatory receptors consisting of CD28, CTLA-4 and PD-1. The interaction of B7-H2/ICOS plays a critical role in Th cell differentiation, T-B cell interactions which is essential for germinal center formation, and humoral immune responses, and as well as the production of cytokine IL-4. In addition, ICOS is more potent in the induction of IL-1 production, a cytokine important for suppressive function of T regulatory cells. The B7-1/B7-2--CD28/CTLA-4 and ICOS-B7RP-1 pathway provides key second signals that can regulate the activation, inhibition and fine-tuning of T-lymphocyte responses. ICOS stimulates both Th1 and Th2 cytokine production but may have a preferential role in Th2 cell development. Moreover, The B7-1/B7-2-CD28/CTLA-4 and ICOS-B7RP-1 pathway has been suggested of being involved in the development of airway inflammation and airway hyperresponsiveness. Cancer Immunotherapy Co-stimulatory Immune Checkpoint Targets Immune Checkpoint Immune Checkpoint Proteins Immune Checkpoint Targets Immunotherapy Targeted Therapy

### Product Info

**Amount :** Mouse ICOS / AILIM / CD278 Recombinant Protein (Fc Tag)(Discontinued) / 200 µg  
**Purification :** > 97 % as determined by SDS-PAGE  
**Content :** Formulation Lyophilized from sterile PBS, pH 7.4  
Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.  
**Storage condition :** Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.  
**Amino Acid :** Met1-Leu142

### Application Note

Endotoxin :< 1.0 EU per µg of the protein as determined by the LAL method




Fig 1: Mouse ICOS / AILIM / CD278 Recombinant Protein (Fc Tag)