

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

37-1182: Mouse c-kit / CD117 Recombinant Protein (Fc Tag)(Discontinued)

Reactivity: Mouse

Bs Protein, Mouse; c-KIT Protein, Mouse; CD117 Protein, Mouse; Fdc Protein, Mouse; Gsfsco1 Protein,

Alternative Name: Mouse; Gsfsco5 Protein, Mouse; Gsfscow3 Protein, Mouse; SCO1 Protein, Mouse; SCO5 Protein, Mouse;

SOW3 Protein, Mouse; Ssm Protein, Mouse; Tr-kit Protein, Mouse; W Protein, Mouse

Description

Source: HEK293 Cells

C-Kit is a type 3 transmembrane receptor for MGF (mast cell growth factor, also known as stem cell factor). c-Kit contains 5 Ig-like C2-type (immunoglobulin-like) domains.and 1 protein kinase domain. It belongs to the protein kinase superfamily, tyr protein kinase family and CSF-1/PDGF receptor subfamily. C-Kit contains 5 Ig-like C2-type (immunoglobulin-like) domains and 1 protein kinase domain. C-Kit has a tyrosine-protein kinase activity. Binding of the ligands leads to the autophosphorylation of KIT and its association with substrates such as phosphatidylinositol 3-kinase. Antibodies to c-Kit are widely used in immunohistochemistry to help distinguish particular types of tumour in histological tissue sections. It is used primarily in the diagnosis of GISTs. In GISTs, c-Kit staining is typically cytoplasmic, with stronger accentuation along the cell membranes. C-Kit antibodies can also be used in the diagnosis of mast cell tumours and in distinguishing seminomas from embryonal carcinomas. Mutations in c-Kit gene are associated with gastrointestinal stromal tumors, mast cell disease, acute myelogenous lukemia, and piebaldism. Defects in KIT are a cause of acute myelogenous leukemia (AML). AML is a malignant disease in which hematopoietic precursors are arrested in an early stage of development. Note=Somatic mutations that lead to constitutive activation of KIT are detected in AML patients. Cancer Immunotherapy Immune Checkpoint Immunotherapy Targeted Therapy

Product Info

Amount: kit / CD117 Recombinant Protein (Fc Tag)(Discontinued) / 100 µg

Purification: > 90 % as determined by SDS-PAGE

Formulation Lyophilized from sterile PBS, pH 7.4

Content: Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before

lyophilization.

Storage condition : Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be

aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Amino Acid: Met1-Thr523

Application Note

Measured by its binding ability in a functional ELISA . Immobilized mouse KITL at 2 $\tilde{A} \square \hat{A} \mu g/ml$ (100 $\tilde{A} \square \hat{A} \mu L/well$) can bind mouse KIT / CD117 with a linear ranger of 1.28-32 ng/ml.

Endotoxin :< 1.0 EU per ̸µg of the protein as determined by the LAL method




Fig 1: Mouse c-kit / CD117 Recombinant Protein (Fc Tag)