

## 37-1186: Mouse CD69 / CLEC2C / AIM Recombinant Protein (Fc Tag, ECD)(Discontinued)

**Reactivity :** Mouse

**Alternative Name :** 5830438K24Rik Protein, Mouse; AI452015 Protein, Mouse; AIM Protein, Mouse; VEA Protein, Mouse

### Description

#### Source : HEK293 Cells

Early activation antigen CD69, also known as activation inducer molecule (AIM), is a single-pass type II membrane protein. Recently, cDNA clones encoding human and mouse CD69 were isolated and showed CD69 to be a member of the C-type lectin superfamily. It is one of the earliest cell surface antigens expressed by T cells following activation. Once expressed, CD69 acts as a costimulatory molecule for T cell activation and proliferation. In addition to mature T cells, CD69 is inducibly expressed by immature thymocytes, B cells, natural killer (NK) cells, monocytes, neutrophils and eosinophils, and is constitutively expressed by mature thymocytes and platelets. CD69 is involved in lymphocyte proliferation and functions as a signal transmitting receptor in lymphocytes, natural killer (NK) cells, and platelets. The structure, chromosomal localization, expression and function of CD69 suggest that it is likely a pleiotropic immune regulator, potentially important in the activation and differentiation of a wide variety of hematopoietic cells. This membrane molecule transiently expresses on activated lymphocytes, and its selective expression in inflammatory infiltrates suggests that it plays a role in the pathogenesis of inflammatory diseases. CD69 plays a crucial role in the pathogenesis of allergen-induced eosinophilic airway inflammation and hyperresponsiveness and that CD69 could be a possible therapeutic target for asthmatic patients.

### Product Info

**Amount :** Mouse CD69 / CLEC2C / AIM Recombinant Protein (Fc Tag, ECD)(Discontinued) / 100 µg

**Purification :** > 90 % as determined by SDS-PAGE.

**Content :** Formulation Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.

**Storage condition :** Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

**Amino Acid :** Gly64-Arg199

### Application Note

Endotoxin :< 1.0 EU per µg protein as determined by the LAL method.




Fig 1: Mouse CD69 / CLEC2C / AIM Recombinant Protein (Fc Tag, ECD)