

37-1213: Human PDGF-C Recombinant Protein (Fc Tag)(Discontinued)

Reactivity : Human
Alternative Name : FALLOTEIN Protein, SCDGF Protein,

Description

Source : HEK293 Cells

PDGF-C is a member of the PDGF/VEGF family of growth factors with a unique domain organization and expression pattern. Platelet-derived growth factor receptors (PDGFRs) are catalytic receptors that have intracellular tyrosine kinase activity. They have roles in the regulation of many biological processes including embryonic development, angiogenesis, cell proliferation and differentiation, and contribute to the pathophysiology of some diseases, including cancer. There are two isoforms of the PDGFR receptor; PDGFRalpha and PDGFRbeta, which can form homo- or heterodimers. The endogenous PDGFR ligands are PDGF-A, -B, -C and -D, which induce receptor dimerization and transphosphorylation at specific tyrosine residues upon binding. This activates the intracellular kinase activity, initiating intracellular signaling through the MAPK, PI 3-K and PKCgamma pathways. PDGF-C acts as a specific ligand for alpha platelet-derived growth factor receptor homodimer, and alpha and beta heterodimer. Binding of this growth factor to its affinity receptor elicits a variety of cellular responses. PDGF-C Appears to be involved in the three stages of wound healing: inflammation, proliferation and remodeling. Involved in fibrotic processes, in which transformation of interstitial fibroblasts into myofibroblasts plus collagen deposition occurs.

Product Info

Amount : C Recombinant Protein (Fc Tag)(Discontinued) / 20 µg
Purification : > 97 % as determined by SDS-PAGE
Content : Formulation Lyophilized from sterile 100mM Glycine, 10mM NaCl, 50mM Tris, pH 7.5
Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.
Storage condition : Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Amino Acid : Val235-Gly345

Application Note

Measured in a cell proliferation assay using Balb/3T3 mouse embryonic fibroblast cells. The ED50 for this effect is typically 0.5-3 µg/mL.
Endotoxin :< 1.0 EU per µg of the protein as determined by the LAL method.
Other pack size also available.




Fig 1: Human PDGF-C Recombinant Protein (Fc Tag)