

37-1265: Human Ephrin-B2 / EFNB2 Recombinant Protein (His & Fc Tag)(Discontinued)

Reactivity : Human

Alternative Name : EPLG5 Protein, Htk-L Protein, HTKL Protein, LERK5 Protein,

Description

Source : HEK293 Cells

EphrinB2 also known as EFNB2 is a member of the ephrin family. EphrinB2 is involved in establishing arterial versus venous identity and perhaps in anastomosing arterial and venous vessels at their junctions. The transmembrane-associated ephrin ligands and their Eph family of receptor tyrosine kinases are expressed by cells of the SVZ. Eph/ephrin interactions are implicated in axon guidance, neural crest cell migration, establishment of segmental boundaries, and formation of angiogenic capillary plexi. Eph receptors and ephrins are divided into two subclasses, A and B, based on binding specificities. Ephrin subclasses are further distinguished by their mode of attachment to the plasma membrane: ephrin-A ligands bind EphA receptors and are anchored to the plasma membrane via a glycosylphosphatidylinositol (GPI) linkage, whereas ephrin-B ligands bind EphB receptors and are anchored via a transmembrane domain. An exception is the EphA4 receptor, which binds both subclasses of ephrins. EphrinB2 expression progressively extends from the arterial endothelium to surrounding smooth muscle cells and to pericytes, suggesting that ephrin-B2 may play an important role during formation of the arterial muscle wall.

Product Info

Amount : B2 / EFNB2 Recombinant Protein (His & Fc Tag)(Discontinued) / 50 µg

Purification : > 97 % as determined by SDS-PAGE

Content : Formulation Lyophilized from sterile PBS, pH 7.4
Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.

Storage condition : Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Amino Acid : Met1-Ala229

Application Note

Measured by its binding ability in a functional ELISA . Immobilized human EphB4-his at 2 µg/ml (100 µL/well) can bind human EphrinB2 Fc chimera with a linear range of 1-25 ng/ml . Immobilized human EphB4 at 2 µg/ml (100 µL/well) can bind human EphrinB2 Fc chimera with a linear range of 1-125 ng/ml.

Endotoxin :< 1.0 EU per µg of the protein as determined by the LAL method.

Other pack size also available.




Fig 1: Human Ephrin-B2 / EFNB2 Recombinant Protein (His & Fc Tag)