

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

37-1271: Human TIMP-1 / TIMP1 Recombinant Protein (Fc Tag)(Discontinued)

Reactivity: Human

Alternative Name: CLGI Protein, EPA Protein, EPO Protein, HCI Protein, TIMP Protein, TIMP-1 Protein,

Description

Source: HEK293 Cells

TIMP metallopeptidase inhibitor 1, also known as TIMP-1/TIMP1, Collagenase inhibitor 16C8 fibroblast Erythroid-potentiating activity, TPA-S1TPA-induced proteinTissue inhibitor of metalloproteinases 1, is a natural inhibitors of the matrix metalloproteinases (MMPs), a group of peptidases involved in degradation of the extracellular matrix. TIMP-1/TIMP1 is found in fetal and adult tissues. Highest levels are found in bone, lung, ovary and uterus. Complexes with metalloproteinases and irreversibly inactivates them by binding to their catalytic zinc cofactor. TIMP-1/TIMP1 mediates erythropoiesis in vitro; but, unlike IL-3, it is species-specific, stimulating the growth and differentiation of only human and murine erythroid progenitors. In addition to its inhibitory role against most of the known MMPs, the protein is able to promote cell proliferation in a wide range of cell types, and may also have an anti-apoptotic function. Transcription of this protein encoding gene is highly inducible in response to many cytokines and hormones. In addition, the expression from some but not all inactive X chromosomes suggests that this gene inactivation is polymorphic in human females. This encoding gene is located within intron 6 of the synapsin I gene and is transcribed in the opposite direction. Complexes with metalloproteinases and irreversibly inactivates them by binding to their catalytic zinc cofactor. TIMP-1/TIMP1 is Known to act on MMP-1, MMP-2, MMP-3, MMP-7, MMP-8, MMP-9, MMP-1, MMP-11, MMP-12, MMP-13 and MMP-16.

Product Info

Amount: 1 / TIMP1 Recombinant Protein (Fc Tag)(Discontinued) / 10 μg

Purification: > 85 % as determined by SDS-PAGE

Formulation Lyophilized from sterile PBS, pH 7.4.

Content : Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before

lyophilization.

Storage condition : Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be

aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Amino Acid: Cys24-Ala207

Application Note

Endotoxin :< 1.0 EU per $\tilde{A} \square \hat{A} \mu g$ protein as determined by the LAL method. Other pack size also available.




Fig 1: Human TIMP-1 / TIMP1 Recombinant Protein (Fc Tag)