

37-1289: Human PTGS2 / COX2 / PGHS-2 Recombinant Protein (His Tag)(Discontinued)

Reactivity : Human
Alternative Name : COX-2 Protein, COX2 Protein, GRIPGHS Protein, hCox-2 Protein, PGG/HS Protein, PGHS-2 Protein, PHS-2 Protein,

Description

Source : Baculovirus-Insect Cells

PTGS2, also known as COX-2, is a component of Prostaglandin-endoperoxide synthase (PTGS). PTGS, also known as cyclooxygenase, is the key enzyme in prostaglandin biosynthesis, and acts both as a dioxygenase and as a peroxidase. There are two isozymes of PTGS: a constitutive PTGS1 and an inducible PTGS2, which differ in their regulation of expression and tissue distribution. PTGS2 is over expressed in many cancers. The overexpression of PTGS2 along with increased angiogenesis and GLUT-1 expression is significantly associated with gallbladder carcinomas. Furthermore the product of COX-2, PGH2 is converted by prostaglandin E2 synthase into PGE2, which in turn can stimulate cancer progression. Consequently inhibiting COX-2 may have benefit in the prevention and treatment of these types of cancer. PTGS2 is regulated by specific stimulatory events, suggesting that it is responsible for the prostanoid biosynthesis involved in inflammation and mitogenesis. It mediates the formation of prostaglandins from arachidonate and may have a role as a major mediator of inflammation and/or a role for prostanoid signaling in activity-dependent plasticity. Cancer Immunotherapy Immune Checkpoint Immunotherapy Targeted Therapy

Product Info

Amount : 2 Recombinant Protein (His Tag)(Discontinued) / 50 µg
Purification : > 90 % as determined by SDS-PAGE
Content : Formulation Lyophilized from sterile 50mM Tris, 100mM NaCl, 0.5mM PMSF, 10% glycerol, pH 8.0
Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.
Storage condition : Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Amino Acid : Met1-Leu604

Application Note

Endotoxin : < 1.0 EU per µg of the protein as determined by the LAL method




Fig 1: Human PTGS2 / COX2 / PGHS-2 Recombinant Protein (His Tag)