

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

37-1329: Human CD45-AB Recombinant Protein (His Tag)(Discontinued)

Reactivity: Human

Alternative Name: B220 Protein, CD45 Protein, CD45R Protein, GP180 Protein, L-CA Protein, LCA Protein, LY5 Protein,

T200 Protein,

Description

Source: HEK293 Cells

The cluster of differentiation (CD) system is commonly used as cell markers in immunophynotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 32 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. Protein tyrosine phosphatase, receptor type C (CD45), also known as PTPRC is a member of the protein tyrosine phosphatase (PTP) family which is known for its function to serve as signaling molecules and to regulate a variety of cellular processes such as cell proliferation, differentiation, mitotic cycle and oncogenic transformation. CD45 is found expression specifically in hemotopietic cells. CD45 consists of an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains. It serves as an essential regulator of T-cell and B-cell antigen receptor signaling through either direct interaction with components of the antigen receptor complexs or by activating various Src family kinases required for the antigen receptor signaling and it also can suppress JAK kinases.

Product Info

Amount: AB Recombinant Protein (His Tag)(Discontinued) / 100 µg

Purification: > 95 % as determined by SDS-PAGE.

Formulation Lyophilized from sterile PBS, pH 7.4.

Content: Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before

lyophilization.

Storage condition : Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be

aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Amino Acid: Met1-Lys529

Application Note

Endotoxin :< 1.0 EU per $\tilde{A} \square \hat{A} \mu g$ protein as determined by the LAL method.




Fig 1: Human CD45-AB Recombinant Protein (His Tag)