

37-1360: Mouse EphA7 / EHK-3 Recombinant Protein (His Tag)(Discontinued)

Reactivity : Mouse
Alternative Name : Cek11 Protein, Mouse; Ebk Protein, Mouse; Ehk3 Protein, Mouse; Hek11 Protein, Mouse; Mdk1 Protein, Mouse; RP23-33D17.1 Protein, Mouse

Description

Source : HEK293 Cells

Ephrin type-A receptor 7, also known as EphA7, belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family which 16 known receptors (14 found in mammals) are involved: EPHA1, EPHA2, EPHA3, EPHA4, EPHA5, EPHA6, EPHA7, EPHA8, EPHA9, EPHA1, EPHB1, EPHB2, EPHB3, EPHB4, EPHB5, EPHB6. The Eph family of receptor tyrosine kinases (comprising EphA and EphB receptors) has been implicated in synapse formation and the regulation of synaptic function and plasticity⁶. Eph receptor-mediated signaling, which is triggered by ephrins⁷, probably modifies the properties of synapses during synaptic activation and remodeling. Ephrin receptors are components of cell signalling pathways involved in animal growth and development, forming the largest sub-family of receptor tyrosine kinases (RTKs). Ligand-mediated activation of Ephs induce various important downstream effects and Eph receptors have been studied for their potential roles in the development of cancer. Down-regulation of EphA7 secondary to hypermethylation has been reported in colorectal cancer. The expression of EphA7 was reduced in all tested gastric cancer cell lines; however, there is marked variability in expression among gastric carcinoma specimens. EphA7 may have roles in the pathogenesis and development of gastric carcinomas.

Product Info

Amount : 3 Recombinant Protein (His Tag)(Discontinued) / 200 µg
Purification : > 97 % as determined by SDS-PAGE
Content : Formulation Lyophilized from sterile PBS, pH 7.4
Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.
Storage condition : Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Amino Acid : Met1-Ile556

Application Note

Measured by its binding ability in a functional ELISA . Immobilized mouse EphA7 at 2 µg/ml (100 µL/well) can bind mouse EphrinA4 with a linear range of 0.08-10 ng/ml .
Endotoxin :< 1.0 EU per µg of the protein as determined by the LAL method




Fig 1: Mouse EphA7 / EHK-3 Recombinant Protein (His Tag)