# **w** abeomics

# 37-1370: Mouse SIRP alpha / CD172a Recombinant Protein (His Tag)(Discontinued)

Reactivity : Mouse

Alternative Name : AI835480 Protein, Mouse; Bit Protein, Mouse; CD172a Protein, Mouse; P84 Protein, Mouse; Ptpns1 Protein, Mouse; SHP-1 Protein, Mouse; SHPS-1 Protein, Mouse; SIRP Protein, Mouse

#### Description

#### Source : HEK293 Cells

Tyrosine-protein phosphatase non-receptor type substrate 1, also known as SHP substrate 1, Inhibitory receptor SHPS-1, Brain Ig-like molecule with tyrosine-based activation motifs, Macrophage fusion receptor, CD172 antigen-like family member A, SIRPA and CD172a, is a single-pass type I membrane protein which contains two Ig-like C1-type (immunoglobulin-like) domains and one Ig-like V-type (immunoglobulin-like) domain. SIRPA is ubiquitously expressed. It is highly expressed in brain and detected at lower levels in heart, placenta, lung, testis, ovary, colon, liver, small intestine, prostate, spleen, kidney, skeletal muscle and pancreas. It is also detected on myeloid cells, but not T-cells. SIRPA is an immunoglobulin-like cell surface receptor for CD47. SIRPA acts as docking protein and induces translocation of PTPN6, PTPN11 and other binding partners from the cytosol to the plasma membrane. SIRPA supports adhesion of cerebellar neurons, neurite outgrowth and glial cell attachment. It may play a key role in intracellular signaling during synaptogenesis and in synaptic function. SIRPA is involved in the negative regulation of receptor tyrosine kinase-coupled cellular responses induced by cell adhesion, growth factors or insulin. It mediates negative regulation of phagocytosis, mast cell activation and dendritic cell activation. Cancer Immunotherapy Co-inhibitory Immune Checkpoint Targets Immune Checkpoint Immune Checkpoint Detection: ELISA Antibodies Immune Checkpoint Detection: FCM Antibodies Immune Checkpoint Immune Checkpoint Targets Immunotherapy Targeted Therapy

## **Product Info**

Amount : Purification :	Mouse SIRP alpha / CD172a Recombinant Protein (His Tag)(Discontinued) / 100 $\mu$ g > 95 % as determined by SDS-PAGE
Content :	Formulation Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.
Storage condition :	Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Amino Acid :	Met1-Asn373

## **Application Note**

Measured by its binding ability in a functional ELISA. Immobilized mouse SIRPA-His at 10  $\tilde{A} \square \hat{A} \mu g/ml$  (100  $\tilde{A} \square \hat{A} \mu L/well$ ) can bind human CD47-Fc , The EC50 of human CD47-Fc is 0.05-0.13  $\tilde{A} \square \hat{A} \mu g/ml$ . Endotoxin :< 1.0 EU per  $\tilde{A} \square \hat{A} \mu g$  of the protein as determined by the LAL method



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com




Fig 1: Mouse SIRP alpha / CD172a Recombinant Protein (His Tag)