

36-1593: Monoclonal Antibody to Beta-2 Microglobulin (Renal Failure & Tumor Marker)(Clone : B2M/961)

Clonality :	Monoclonal
Clone Name :	B2M/961
Application :	FACS,IF,WB,IHC
Reactivity :	Human
Gene :	B2M
Gene ID :	567
Uniprot ID :	P61769
Format :	Purified
Alternative Name :	B2M,CDABP0092,HDCMA22P
Isotype :	Mouse IgG2b, kappa
Immunogen Information :	Full length recombinant human B2M protein

Description


Recognizes a protein of 12kDa, identified as β -2 microglobulin. Major histocompatibility complex (MHC) class 1 molecules bind to antigens for presentation on the surface of cells. The proteasome is responsible for producing these antigens from the components of foreign pathogens. MHC class 1 molecules consist of an α heavy chain that contains three subdomains (α 1, α 2, α 3) and a non-covalent associating light chain, known as β -2-Microglobulin. β -2-Microglobulin associates with the α 3 subdomain of the α heavy chain and forms an immunoglobulin domain-like structure that mediates proper folding and expression of MHC class 1 molecules. The α 1 and α 2 domains of the α heavy chain form the peptide antigen-binding cleft. Mutations in the β -2-Microglobulin gene can enhance the progression of malignant melanoma phenotypes.

Product Info


Amount :	100 μ g
Purification :	Affinity Chromatography
Content :	100 μ g in 500 μ l PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
Storage condition :	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note


Flow Cytometry (0.5-1 μ g/million cells); ,Immunofluorescence (1-4 μ g/ml); Western Blot (0.5-2 μ g/ml);,Immunohistochemistry (Formalin-fixed) (0.5-1 μ g/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 $\text{\textcircled{C}}$ followed by cooling at RT for 20 minutes),




Formalin-fixed, paraffin-embedded human Cervical Carcinoma stained with Beta-2-Microglobulin Monoclonal Antibody (B2M/961).




Formalin-fixed, paraffin-embedded human Melanoma stained with Beta-2-Microglobulin Monoclonal Antibody (B2M/961).




Formalin-fixed, paraffin-embedded human Endometrial Carcinoma stained with Beta-2-Microglobulin Monoclonal Antibody (B2M/961)




Formalin-fixed, paraffin-embedded human Renal Carcinoma stained with Beta-2-Microglobulin Monoclonal Antibody (B2M/961)



Formalin-fixed, paraffin-embedded human Cervical Carcinoma stained with Beta-2-Microglobulin Monoclonal Antibody (B2M/961)



Formalin-fixed, paraffin-embedded human Bladder Carcinoma stained with Beta-2-Microglobulin Monoclonal Antibody (B2M/961)



Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with Beta-2-Microglobulin Monoclonal Antibody (B2M/961)