

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-2027: Anti-p27Kip1 (Mitotic Inhibitor/Suppressor Protein) Monoclonal Antibody (Clone: KIP1/769)-CF488

Clonality: Monoclonal **Clone Name:** KIP1/769 Application: FACS,IF

Reactivity: Human, Mouse, Rat

Conjugate: CF488 Gene: CDKN1B Gene ID: 1027 **Uniprot ID:** P46527

CDKN1B, CDKN4, Cyclin Dependent Kinase Inhibitor 1B, Cyclin-dependent kinase inhibitor p27 **Alternative Name:**

Kip1, KIP1, MEN1B, MEN4

Isotype: Mouse IgG1, kappa

Immunogen Information: Recombinant human KIP1 protein

Description

This MAb recognizes a 27kDa protein, identified as the p27Kip1, a cell cycle regulatory mitotic inhibitor. It is highly specific and shows no cross-reaction with other related mitotic inhibitors. In Western blotting of cell lysates from 7 human breast cancer cell lines (ZR75-1, ZR75-30, MCF-7, MDAMB453, T47D, CAL51, 734B), the antibody labels a single band corresponding to p27Kip1. It functions as a negative regulator of G1 progression and has been proposed to function as a possible mediator of TGF- induced G1 arrest. p27Kip1 is a candidate tumor suppressor gene. Reportedly, low p27 expression has been associated with unfavorable prognosis in renal cell carcinoma, colon carcinoma, breast carcinomas, non-small-cell lung carcinoma, hepatocellular carcinoma, multiple myeloma, and lymph node metastases in papillary carcinoma of the thyroid, as well as a more aggressive phenotype in carcinoma of the cervix.

Product Info

Amount: 0.5 ml at 100µg/ml

Antibody Purified from Bioreactor Concentrate by Protein A/G and conjugated to various

Content: reporter molecules. Prepared in 10mM PBS with 0.05% BSA and 0.05% azide. Contact us if you

require this Ab in a different format.

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody Storage condition:

is stable for 24 months.

Application Note

Flow Cytometry (5ul per test per one million cells or 5ul per 100ul of whole blood); Immunofluorescence (1:50-1:100);




Fig.1: Flow Cytometry of human p27 on Jurkat cells. Black: cells alone; Grey: Isotype Control; Green: CF488-labeled p27 Monoclonal Antibody (KIP1/769).