

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 36-2040: Anti-Carcinoembryonic Antigen (CEA) / CD66 Monoclonal Antibody (Clone: C66/1009)-CF488

Clonality: Monoclonal
Clone Name: C66/1009
Application: FACS,IF
Reactivity: Human
Conjugate: CF488

Gene: CEACAM1 & CEACAM5

**Gene ID:** 1048; 634 **Uniprot ID:** P06731

Alternative Name: Carcinoembryonic Antigen-related Cell Adhesion Molecule 5, CEACAM5, CD66, Biliary

Glycoprotein (BGP-1)

**Isotype:** Mouse IgG2a, kappa

Immunogen Information: Recombinant full-length human CEA protein

## **Description**

This antibody recognizes proteins of 80-200kDa, identified as different members of CEA family. CEA is synthesized during development in the fetal gut and is re-expressed in increased amounts in intestinal carcinomas and several other tumors. This MAb does not react with nonspecific cross-reacting antigen (NCA) and with human polymorphonuclear leucocytes. It shows no reaction with a variety of normal tissues and is suitable for staining of formalin/paraffin tissues. CEA is not found in benign glands, stroma, or malignant prostatic cells. Antibody to CEA is useful in detecting early foci of gastric carcinoma and in distinguishing pulmonary adenocarcinomas (60-70% are CEA+) from pleural mesotheliomas (rarely or weakly CEA+). Anti-CEA positivity is seen in adenocarcinomas from the lung, colon, stomach, esophagus, pancreas, gallbadder, urachus, salivary gland, ovary, and endocervix.

## **Product Info**

Amount: 0.5 ml at 100µg/ml

Antibody Purified from Bioreactor Concentrate by Protein A/G and conjugated to various

**Content:** reporter molecules. Prepared in 10mM PBS with 0.05% BSA and 0.05% azide. Contact us if you

require this Ab in a different format.

**Storage condition :** Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months.

## **Application Note**

Flow Cytometry (5ul per test per one million cells or 5ul per 100ul of whole blood); Immunofluorescence (1:50-1:100);